For patients and caregivers

Alkindi Sprinkle® (hydrocortisone) oral granules

Even slightly off... is not good enough.

In pediatric adrenal insufficiency, getting the dose right matters

ALKINDI SPRINKLE® (hydrocortisone) oral granules: The first hydrocortisone adrenal insufficiency treatment that is specifically designed to help provide accurate and individualized dosing for newborns and children.

USE

ALKINDI SPRINKLE is a prescription medicine used in children from birth to less than 17 years old as a replacement therapy when the adrenal gland is not making enough cortisol.

IMPORTANT SAFETY INFORMATION

Always give ALKINDI SPRINKLE exactly as your doctor has directed.

Do not take ALKINDI SPRINKLE if you are allergic to hydrocortisone or any of its other ingredients.

Please see additional Important Safety Information throughout and enclosed full Prescribing Information.

Even slightly off is not good enough. Too much or too little hydrocortisone can have lasting consequences

Dosing accuracy in children with adrenal insufficiency isn't optional—it's essential. If your child gets too much or too little hydrocortisone, the consequences can last for years, even into adulthood.

Too much hydrocortisone can cause long-term harm

If your child consistently receives **more hydrocortisone** than prescribed, they are at risk for serious long-term health problems that can follow them into adulthood.

- Growth failure
- Cushing's syndrome
- Abnormal weight gain
- High blood pressure

Too little hydrocortisone can put your child in immediate danger

If your child gets **less hydrocortisone** than prescribed, they could be in immediate danger.

- Adrenal crisis
- Rapid growth
- Weight loss
- Masculine features
- Poor response to stress

No representation is made regarding the use of ALKINDI SPRINKLE® (hydrocortisone) oral granules in relation to these conditions.

ALKINDI SPRINKLE: The first FDA-approved hydrocortisone designed specifically for newborns and children.

IMPORTANT SAFETY INFORMATION

Adrenal Crisis: giving too low a dose or stopping medication can cause low levels of cortisol, which can result in serious illness or death. Treatment with intravenous hydrocortisone should be started immediately. When switching from another type of hydrocortisone to ALKINDI SPRINKLE, watch your child closely for any changes. If your child doesn't get the entire dose of ALKINDI SPRINKLE because of vomiting or spitting some granules out, contact your doctor to see if another dose is needed.

Alkindi Sprinkle®
(hydrocortisone)
oral granules

The dose your doctor prescribed. Designed for accurate delivery, every time

ALKINDI SPRINKLE is a granular version of hydrocortisone, which means it's small enough to sprinkle onto soft food or directly into your child's mouth. ALKINDI SPRINKLE capsules are designed to be opened and then the granules are given on either a full or empty stomach with a sip of fluids. ALKINDI SPRINKLE has a gritty texture that may be new to most children.

Here are some useful suggestions on how to give ALKINDI SPRINKLE to your child.

ON A SPOON WITH FOOD

Pour **all** granules onto a spoonful of soft food, like yogurt or applesauce, and give within 5 minutes.

DIRECTLY INTO YOUR CHILD'S MOUTH

Pour **all** granules of the prescribed dose directly into your child's mouth.



Scan this code for access to useful videos for giving ALKINDI SPRINKLE to your child.

IMPORTANT SAFETY INFORMATION

Immunosuppression and Increased Risk of Infection with Use of a Dosage Greater Than Replacement: Use of a greater than replacement dosage can suppress the immune system and increase the risks of new infections or exacerbation of latent infections with any pathogen, including viral, bacterial, fungal, protozoan, or helminthic infections. Monitor patients for signs and symptoms of infections.

Important Safety Information continued on next page.

Please see additional Important Safety Information throughout and enclosed full Prescribing Information.

ALKINDI SPRINKLE: The FDA-approved hydrocortisone formulation specifically designed for newborns and children—delivering precise, premeasured, low-dose strengths to help with accurate pediatric dosing

ALKINDI SPRINKLE® (hydrocortisone) oral granules are provided in color-coded capsules for easy identification and to help with accurate dosing.

- Available in 4 flexible dosage strengths (0.5 mg, 1 mg, 2 mg, and 5 mg)
- Granules are less than 0.8 mm in size, much smaller than the FDA-recommended size to prevent choking
- Stored in easy-to-open capsules

ALKINDI SPRINKLE can be **individualized to your child's specific cortisol need with the lowest possible dose used**.

4 flexible strengths to assist accurate dosing



Capsules shown are not actual size.

IMPORTANT SAFETY INFORMATION

Growth Retardation: the long-term use of corticosteroids in high doses may cause growth retardation in children.

Decrease in Bone Density: corticosteroids can affect your child's bone growth and strength.

Cushing's Syndrome Due to High Doses of Corticosteroids: treatment with high doses of corticosteroids can cause Cushing's Syndrome. Treatment should be limited to the smallest dose required, and your child's growth and development monitored appropriately.

Alkindi Sprinkle® (hydrocortisone) oral granules

ALKINDI SPRINKLE: No pill splitting. No pill crushing. Just designed for accurate dosing every time[1]*

1 Check expiration and prepare

Check the expiration date on the ALKINDI SPRINKLE bottle. **Do not** use after the expiration date listed on the bottle.

Then, remove the prescribed dose of ALKINDI SPRINKLE capsules from the bottle.

Capsules shown are not actual size.

2 Hold and tap

Hold capsule with the writing at the top. Tap the capsule to make sure the granules fall to the bottom.

3 Squeeze

Gently squeeze the bottom of the capsule to loosen the top from the bottom.

4 Twist and remove

Carefully twist off the top of the capsule.

5 Sprinkle

ALKINDI SPRINKLE can be given by spoon with soft food or sprinkled directly into the child's mouth.

6 Tap

Tap the capsule to ensure all the granules are removed. Avoid wetting the capsule on the tongue or soft food as this may result in granules remaining in the capsule.

7 Give fluids

After giving ALKINDI SPRINKLE, give the child a sip of fluids, such as water, milk, breast milk, or formula, right away to ensure all granules are swallowed.

*When the entire dose is given as directed.

ALKINDI SPRINKLE has a texture that may be new to most children

Some pediatric patients may object to the texture of ALKINDI SPRINKLE in their mouth. It's best to give with soft food. ALKINDI SPRINKLE can also be poured directly into the child's mouth.

ALKINDI SPRINKLE granules may sometimes be seen in stools because the center of the granule is not absorbed in the gut after the drug has been released

IMPORTANT SAFETY INFORMATION

Changes in Vision: tell your doctor if your child has blurred vision or other vision problems during treatment with ALKINDI SPRINKLE.

Important Safety Information continued on next page.

Please see additional Important Safety Information throughout and enclosed full Prescribing Information.

Your rare disease journey is personal. So is our support

When your doctor prescribes ALKINDI SPRINKLE® (hydrocortisone) oral granules, you're automatically enrolled in the Eton Cares Program®*—a full-service support program designed to help you and your child at every step of the treatment journey.

Nurse ambassador support

- Regular check-ins by phone
- Answer health or medication questions
- Help you and your child stay on track with treatment

Pharmacists

- 24/7 support
- Refill reminders
- Shipment alerts

Insurance specialists

- Benefits verification and prior authorization assistance
- Help securing financial support through the Eton Cares Patient Assistance Program*
- Referrals to other sources of financial assistance if needed

When your doctor sends your prescription to our specialty pharmacy, Anovo®, here is what you can expect:

1 | **Welcome call** from your Care Coordinator to verify your prescription, offer support, and answer questions

2 | **Insurance verification** to review coverage and benefits

3 | **Financial assistance review** to explore copay programs and other support options

4 | **Medication delivery** with instructions and product-specific information

5 | **Ongoing pharmacy support**, including refill coordination, reminders, and 24/7 access to a pharmacist

6 | **Nurse ambassador support** to help you and your child stay on track with treatment, answer health or medication questions, and provide regular check-ins by phone

*Restrictions, limitations, and/or eligibility requirements may apply.

IMPORTANT SAFETY INFORMATION

Psychiatric Changes: corticosteroids can change your child's behavior or mood. Tell your doctor if your child has periods of extreme happiness, extreme sadness, hallucinations, or depression.

Please see additional Important Safety Information on the next page.

Alkindi Sprinkle® (hydrocortisone) oral granules

$0 copay for eligible patients*

For questions or support, please contact us at our specialty pharmacy, Anovo

Call 1-833-343-2500 Monday through Friday, 8:00 AM-5:00 PM CT. Anovo's Pharmacy Team is always available 24 hours a day, 365 days a year.

A helpful tip: We recommend saving Anovo's phone number in your contacts so you can easily recognize important calls about your prescription, refills, or insurance updates.

*Commercially eligible patients can pay $0 per month. Restrictions, limitations, and/or eligibility requirements may apply.

IMPORTANT SAFETY INFORMATION

Gastrointestinal Reactions: tell the doctor if your child has stomach pain, upset stomach, black, tarry stools, or vomiting of blood. These could be signs of ulcers or tears in the stomach or intestines. Taking anti-inflammatory nonsteroidal drugs, like ibuprofen, naproxen, or aspirin, can increase the risk of ulcers or tears.

The most common side effects of ALKINDI SPRINKLE include retaining fluids, changes in glucose tolerance, high blood pressure, behavioral and mood changes, greater appetite, and weight gain.

Vaccination: Administration of live vaccines may be acceptable in ALKINDI SPRINKLE-treated pediatric patients with adrenocortical insufficiency who receive replacement corticosteroids.

Please see additional Important Safety Information throughout and enclosed full Prescribing Information.

For patients and caregivers

Alkindi Sprinkle® (hydrocortisone) oral granules

Don't accept imprecision. Get started with ALKINDI SPRINKLE

The adrenal insufficiency treatment designed specifically for newborns and children

Learn more at ALKINDISPRINKLE.com.

USE AND IMPORTANT SAFETY INFORMATION

USE

ALKINDI SPRINKLE® (hydrocortisone) oral granules is a prescription medicine used in children from birth to less than 17 years old as replacement therapy when the adrenal gland is not making enough cortisol.

IMPORTANT SAFETY INFORMATION

Always give ALKINDI SPRINKLE exactly as your doctor has directed.
Do not take ALKINDI SPRINKLE if you are allergic to hydrocortisone or any of its other ingredients.

Adrenal Crisis: giving too low a dose or stopping medication can cause low levels of cortisol, which can result in serious illness or death. Treatment with intravenous hydrocortisone should be started immediately. When switching from another type of hydrocortisone to ALKINDI SPRINKLE, watch your child closely for any changes. If your child doesn't get the entire dose of ALKINDI SPRINKLE because of vomiting or spitting some granules out, contact your doctor to see if another dose is needed.

Immunosuppression and Increased Risk of Infection with Use of a Dosage Greater Than Replacement: Use of a greater than replacement dosage can suppress the immune system and increase the risks of new infections or exacerbation of latent infections with any pathogen, including viral, bacterial, fungal, protozoan, or helminthic infections. Monitor patients for signs and symptoms of infections.

Growth Retardation: the long-term use of corticosteroids in high doses may cause growth retardation in children.

Decrease in Bone Density: corticosteroids can affect your child's bone growth and strength.

Cushing's Syndrome Due to High Doses of Corticosteroids: treatment with high doses of corticosteroids can cause Cushing's Syndrome. Treatment should be limited to the smallest dose required, and your child's growth and development monitored appropriately.

Changes in Vision: tell your doctor if your child has blurred vision or other vision problems during treatment with ALKINDI SPRINKLE.

Psychiatric Changes: corticosteroids can change your child's behavior or mood. Tell your doctor if your child has periods of extreme happiness, extreme sadness, hallucinations, or depression.

Gastrointestinal Reactions: tell the doctor if your child has stomach pain, upset stomach, black, tarry stools, or vomiting of blood. These could be signs of ulcers or tears in the stomach or intestines. Taking anti-inflammatory nonsteroidal drugs, like ibuprofen, naproxen, or aspirin, can increase the risk of ulcers or tears.

The most common side effects of ALKINDI SPRINKLE include retaining fluids, changes in glucose tolerance, high blood pressure, behavioral and mood changes, greater appetite, and weight gain.

Vaccination: Administration of live vaccines may be acceptable in ALKINDI SPRINKLE-treated pediatric patients with adrenocortical insufficiency who receive replacement corticosteroids.

Please visit ALKINDISPRINKLE.com/patient for more information.

You are encouraged to report negative side effects of prescription drugs by contacting Eton Pharmaceuticals, Inc. at 1-855-224-0233 or the U.S. Food and Drug Administration (FDA) at www.fda.gov/safety/medwatch or call 1-800-FDA-1088.

Please see enclosed full Prescribing Information for more information.

eton PHARMACEUTICALS

© 2026 Eton Pharmaceuticals. All rights reserved. Eton, ALKINDI SPRINKLE, and Eton Cares are trademarks of Eton Pharmaceuticals. All other trademarks are property of their respective owners.
1569-v3